

On multiple connectedness of regions visible due to multiple diffuse reflections

Sudebkumar Prasant Pal¹

Department of Computer Science and Engineering,
Indian Institute of Technology, Kharagpur

Kharagpur 721302, India.

email: spp@cse.iitkgp.ernet.in

url:- <http://www.angelfire.com/or/sudebkumar>

Dilip Sarkar

Department of Computer Science
University of Miami, Coral Gables

Miami, FL 33124, USA

email: sarkar@cs.miami.edu

Abstract

It is known that the region $V(s)$ of a simple polygon P , directly visible (illuminable) from an internal point s , is simply connected. Aronov et al. [2] established that the region $V_1(s)$ of a simple polygon visible from an internal point s due to at most one diffuse reflection on the boundary of the polygon P , is also simply connected. In this paper we establish that the region $V_2(s)$, visible from s due to at most two diffuse reflections may be multiply connected; we demonstrate the construction of an n -sided simple polygon with a point s inside it so that and the region of P visible from s after at most two diffuse reflections is multiple connected.

Keywords: multiple connectedness, diffuse reflection, visibility, simple polygon.

¹A part of this work was done when this author was visiting the University of Miami, Coral Gables, Florida, USA.

1 Introduction

In recent years, visibility problems have been studied extensively (see [8, 11]). Visibility computations abound in computer graphics, motion planning, robotics and computer vision. Two points inside a simple polygon are mutually *visible* if the line segment joining them is not obstructed by any edge of the polygon. Several algorithms exist for computing the region visible from a point light source inside a simple polygon [4, 6, 5]. The problems of computing the region of a simple polygon which is *weakly visible* from an internal segment [7, 5], or a convex set [3], are also well studied. A point p inside a simple polygon P is said to be *weakly visible* from an edge e of the polygon if p is visible from some point in the *interior* of e . Certain portions of the polygon that are not directly illuminated from the source may become visible due to one or more reflections on the bounding edges of the polygon. Reflection at a point is called *specular* if the reflected ray is directed as per the standard law of reflection; the angle of incidence is the same as the angle of reflection. Most reflecting surfaces cause another type of reflection, called *diffuse* reflection, where light incident at a point is reflected in all possible interior directions. We assume that light incident at a vertex is absorbed and not reflected.

Throughout this paper we consider the region visible from a point source s due to multiple reflections inside a simple polygon P . Such visible regions are also called *visibility polygons*.

1.1 Preliminaries and notation

Let P be a simple polygon with no three collinear vertices. Let $int(X)$ and $bd(X)$ denote the relative interior and boundary of a region X of P , respectively. If p and q are two points on $bd(P)$, let $bd(p, q)$ denote the part of $bd(P)$ traversed from p to q in counterclockwise order, keeping $int(P)$ to the left of the direction of traversal. Two points in P are (*mutually*) *visible* if the interior of the line segment joining them lies in $int(P)$. A point y is *visible under diffuse reflection* from a point x , if there exists a point p lying in the interior of an edge of P , such that p is visible from both x and y .




Figure 1.1: Visibility with multiple diffuse reflections.

Let S denote a point light source inside P . A point y is 1-visible from S if y and S are mutually visible. For $k > 1$, a point $y(= p_k)$ is k -visible from a point $S(= p_0)$, if there exists

points $p_1, p_2, \dots, p_{k-1}$ in the interiors of edges of P , such that p_i and p_{i+1} are mutually visible for $0 \leq i \leq k-1$. We say that the ray emitted from S reaches y after $k-1$ *diffuse reflections* at points $p_1, p_2, \dots, p_{k-1}$. In Figure 1.1, y is *2-visible* from S ; y is visible from S after one diffuse reflection. We assume that the light incident at a vertex is absorbed and not reflected further.

Let $V(s)$ denote the portion of P visible from s . We know that $V(s)$ is simply connected, has $O(n)$ edges, and at most one edge of $V(s)$ lies on each edge of P [4, 5]. For a point $S \in P$, let $V(S) = V(P, S)$ denote the polygonal region consisting of points in P that are directly visible from S . For $k \geq 0$, let $V_k(S)$ denote the polygonal region consisting of points that are l -visible from S , for some $1 \leq l \leq k+1$. $V_k(s)$ is the set of points that receive light from s after at most k diffuse reflections. We have $V_0(s) = V(s)$.

1.2 Combinatorial complexity and multiple connectedness of regions visible due to diffuse reflections

Regions visible due to reflections inside simple polygons were first studied by Aronov *et al.* [2]; a tight $\Theta(n^2)$ bound was established on the number of edges of visibility polygons when at most one diffuse reflection is permitted. The authors showed that such visibility polygons were simply connected. In sharp contrast, regions visible due to at most one specular reflection were shown to be multiply connected, with combinatorial complexity $\Theta(n^2)$ [2]. (The combinatorial complexity of a visibility polygon is the number of edges defining its boundary.) Indeed, the number of *holes* (also called *blind spots*) in such a visibility polygon (with at most one specular reflection) is $\Theta(n^2)$ [2].

The case where at most k specular reflections are permitted, is dealt with in a subsequent paper [1]; an $O(n^{2k})$ upper bound and a lower bound of $\Omega((n/k - \Theta(1))^{2k})$ on the combinatorial complexity of the visibility polygons was established when at most a constant number k of specular reflections are permitted. The number of holes too obeys these bounds. The upper bound on the combinatorial complexity of visibility polygons improves to $O(n^{2\lceil(k+1)/2\rceil+1})$, when specular reflections are replaced by diffuse reflections [10]. Note that the improvement is by an asymptotic square root factor for large k . This is not surprising because diffuse reflection spreads light much more widely than restricted pencils do in the case of specular reflections.

The best lower bound on the combinatorial complexity for visibility polygons $V_k(s)$ (with at most k diffuse reflections), for any $k \geq 1$, remains just $\Omega(n^2)$ to date (see [2, 10]). One interesting and important problem concerning multiple diffuse reflections is that of bridging this gap between the very high upper bound of [10] and the $\Omega(n^2)$ lower bound of [2, 10]. One way to approach this open problem is to study the structure of $V_k(s)$ and bound the number of holes in it. Showing that the number of holes is not as high as in the case of specular reflections (see [1]), may help improving the upper bound. The presence of holes in $V_2(s)$ (and

in $V_k(s)$ for $k \geq 1$, in general) was not known until we demonstrated the existence of holes in $V_k(s)$, $k \geq 1$, in [9]. This paper presents essentially the construction in [9] of an n -sided simple polygon P with a point s located inside P so that $V_2(s)$ has a hole. We construct a simple polygon P and place a point s inside it so that the portion of the polygon P visible from s due to at most two diffuse reflections does not include the interior of a triangle tqr (see Figures 2.2 and 2.3). Here, the triangle tqr is totally contained inside P and the boundary of triangle tqr as well as its exterior neighbourhood are visible from s after at most two diffuse reflections.

An interesting problem is to find asymptotic bounds on the number of holes in $V_2(s)$. We believe that there can not be more than a linear number of holes. Such a linear bound on the number of holes in $V_k(s)$ for any $k \geq 1$ may improve the upper bound on the combinatorial complexity of $V_k(s)$ over that in [10].

The existence of holes in $V_2(s)$ is a bit surprising. Diffuse reflection, as the term suggests, results in bouncing back of incident light in all possible directions from the point of reflection. Clearly, a lot more region gets illuminated due to such reflections when compared with specular reflection. Multiple diffuse reflections would lead to even larger spreads and it was not intuitively apparent as to how holes could be created in such widely spread visible regions. In fact, it seems unlikely that too many such holes can be created; we suspect that the number of holes in $V_2(s)$ is $O(n)$.

2 The construction of a hole for two diffuse reflections

Now we show that $V_2(s)$ may have a hole. In what follows, we state this result in detail as in [9]. We show the existence of an island or hole by constructing a simple polygon P containing a point s such that $V_2(s)$ has a triangular hole with vertices t , q and r (see Figures 2.2 and 2.3). Note that rays emanating from s are drawn with dashed lines. Rays that emerge after the first reflection (from points visible to s) are drawn with normal lines. Rays that emerge from points of $bd(V_1(s))$ (after the second reflection) are drawn with lines having alternating dots and dashes. The interior of the triangle tqr is invisible from s , from all points of $bd(V_0(s))$, and, from all points on $bd(V_1(s))$. Note that the first reflection takes place on points of $bd(V_0(s))$ and the second reflection takes place on points of $bd(V_1(s))$.

We need a definition. We say that a point p *blocks* a region R of P from another region S if the shortest path from every point in R to every point in S passes through p .

Clearly, no ray can reach the hole directly from s . This is because b blocks s from the (proposed) hole.

LEMMA 2.1 The triangle tqr is not visible from s after the first reflection.

PROOF.




Figure 2.2 Regions visible directly and after one reflection.




Figure 2.3. Region visible due to one reflection.

Consider $bd(V_0(s))$. We start by classifying portions of $bd(P)$ that form $bd(V_0(s))$. There are three portions- $bd(x, y)$, $bd(w, b)$ and $bd(a, v)$. These constitute all points on $bd(V_0(s))$. The sequences of points, $\langle s, a, y \rangle$, $\langle s, b, x \rangle$ and $\langle s, v, w \rangle$ are collinear. How do we guard the hole from the points of $bd(V_0(s))$? The shortest path from any point in $bd(x, y)$ to any point in the hole has to pass through j . In other words, j blocks the hole from $bd(x, y)$. So, no point in $bd(x, y)$ can see the hole. Similarly, b blocks $bd(w, b)$ from the holes, and, c blocks $bd(a, v)$ from the hole. So, no point on $bd(V_0(s))$, can see the hole. We conclude that the hole is not visible from any point of $bd(V_0(s))$.

□

All we need to do now is to show that $bd(V_1(s))$ can not see the hole. We start by classifying portions of $bd(P)$ that form $bd(V_1(s))$. These are

1. Points of $bd(a, b)$ visible from $bd(a, v)$, $bd(w, b)$ or $bd(x, y)$. (Note that $bd(x, y)$ is the only portion of $bd(P)$ visible from s outside $bd(a, b)$.)
2. Points of $bd(i, f)$ and $bd(b, c)$, visible from $bd(a, v)$. (Points of $bd(a, v)$ can see points within $bd(i, f)$ because (i) the vertex i is to the right of the ray starting at a and passing through c , and, (ii) the vertex f is to the left of the ray starting at v and passing through a . Observe that $bd(a, v)$ can also see points on $bd(b, c)$. These are the only portions of $bd(P)$, visible from $bd(a, v)$.)
3. Points of $bd(h, a)$ visible from $bd(w, b)$. (Points in $bd(w, b)$ can see points of $bd(P)$ only within $bd(d, a)$, or more precisely, points within $bd(d, k)$, where e, b, a and k are collinear.)
4. Points of $bd(b, c)$ visible from $bd(x, y)$. (The point c is to the left of $ray(y, j)$. So, all points of $bd(V_1(s))$ on $bd(b, j)$ are to the right of $ray(y, j)$).
5. Points of $bd(h, a)$ visible from $bd(x, y)$.
6. Points of $bd(i, h)$ visible from $bd(x, y)$.

The remaining points on $bd(P)$ can not be on $bd(V_1(s))$.

Now we show that none of these six possible portions of $bd(V_1(s))$ can illuminate the triangular hole tqr of $V_2(s)$.

1. The second reflection at $bd(a, b)$ causes rays to miss the hole due to the blocker c . This is due to the fact that the shortest path from any point in $bd(a, b)$ to any point in the hole tqr , has to cross segment ac . In particular, points in $bd(v, b)$ are blocked from the hole by b .

2. Consider points of $bd(V_1(s))$ on $bd(i, f)$. It is sufficient to consider the extreme points y' and c' of $bd(i, f)$ in $bd(V_1(s))$. The point y' is collinear to v and a . The point j blocks y' and the whole of $bd(j, y')$ from the hole. The point c' is collinear with a and c . The point i blocks the whole of $bd(i, j)$ from the hole tqr .

Now consider points of $bd(b, c)$ that belong to $bd(V_1(s))$ due to first reflection at $bd(a, v)$. These points are blocked from the hole by c .

3. Only points within $bd(d, k)$ in $bd(h, a)$ can cause the second reflection from rays undergoing the first reflection on $bd(w, b)$. All these points in $bd(d, k)$ are blocked by d from the hole tqr . Indeed, k, d, q and r are collinear, defining the edge qr of tqr . Note that q is the intersection of $ray(k, d)$ and $ray(f, i)$ and, r is the intersection of $ray(k, d)$ and $ray(g, h)$.
4. Consider points of $bd(b, c)$ visible from xy . Points of $bd(b, c)$ can not see tqr because they are blocked by c .
5. Consider $bd(d, a)$. Points in $bd(d, z)$ are blocked by h from points on xy . Since z, h and x are collinear and tqr is to the right of $ray(z, c)$, all points of $bd(z, a)$ are blocked from tqr by c .
6. Consider $bd(i, h)$ and xy . Points on $bd(j, y)$ are blocked from tqr by j . So we need to consider only $bd(y, h)$. Any point on $bd(y, f)$ may reflect into the region to the right of $ray(f, i)$ but not to the left of $ray(f, i)$. In the construction, f, i, t and q are collinear and tq defines an edge of the hole tqr . Note that $ray(g, h)$ and $ray(f, i)$ intersect at vertex t .

Now consider point f' on $bd(f, h)$ where f' is the point where $ray(x, f)$ meets $bd(P)$. We construct $bd(f, g)$ so that the shortest path from f' to h passes through g and g lies in the segment joining f and f' . Points on xy can not see any point in the interior of $bd(f, f')$. Since g, h, t and r are collinear, (i) the shortest path from any point in $bd(f', g)$ to any point inside tqr has to pass through blockers h and g , and, (ii) the shortest path from any point in $bd(g, h)$ to any point in tqr has to pass through blocker h . Indeed, $ray(g, h)$ defines the edge tr of hole tqr .

Note that reflections on edge $g'h$ will not reach the interior of triangle tqr because triangle tqr is to the right of $ray(g', h)$.

In summary, we have established that tqr is a hole in $V_2(s)$. We state the result as a theorem.

THEOREM 2.2 It is possible to construct a simple polygon P with internal points s, t, q and r such that (i) tqr is a triangle totally contained in $int(P)$ (ii) all points inside the triangle tqr belong to $P \setminus V_2(s)$ and (iii) tq, tr and rq are on $bd(V_2(s))$.

3 Concluding remarks

We have shown that the region visible due to at most two diffuse reflections can have holes. This result holds for single point light sources as well as single extended light sources like a line segment or a circle. The details are simple and are omitted here. For three or more diffuse reflections too, such multiply connected visible regions can be created by extending our construction; the source s in our construction can be created as an illuminating line segment on an edge of the polygon after a predetermined number $k - 2$, $k \geq 3$ of reflections from a point s' , where light from s' must undergo $k - 2$ reflections (inside say, a winding spiral) and hit s before spreading to the rest of the polygon. The details are straightforward and are omitted here.

An important research direction is that of establishing upper bounds on the number of holes in diffuse visibility regions. We believe that the upper bound is as low as linear in the number of vertices of the input polygon.

References

- [1] B. Aronov, A. Davis, T. K. Dey, S. P. Pal, and D. C. Prasad. Visibility with multiple reflections. *Discrete and Computational Geometry*, 20:61–78, 1998.
- [2] B. Aronov, A. Davis, T. K. Dey, S. P. Pal, and D. C. Prasad. Visibility with one reflection. *Discrete and Computational Geometry*, 19:553–574, 1998.
- [3] S. K. Ghosh. Computing the visibility polygon from a convex set and related problems. *Journal of Algorithms*, 12:75–95, 1991.
- [4] H. El Gindy and D. Avis. A linear algorithm for computing the visibility polygon from a point. *Journal of Algorithms*, 2:186–197, 1981.
- [5] L. Guibas, J. Hershberger, D. Leven, M Sharir, and R. Tarjan. Linear time algorithms for visibility and shortest path problems inside triangulated simple polygons. *Algorithmica*, 2:209–233, 1987.
- [6] D. T. Lee. Visibility of a simple polygon. *Computer Vision, Graphics and Image Processing*, 22:207–221, 1983.
- [7] D. T. Lee and A. K. Lin. Computing the visibility polygon from an edge. *Computer Vision, Graphics and Image Processing*, 34:1–19, 1986.
- [8] J. O’Rourke. *Art gallery theorems and algorithms*. Oxford University Press, 1987.

- [9] S. P. Pal and D. Sarkar. On multiple-connectedness of regions visible due to multiple diffuse reflections. *Technical Report No. TR/IIT/CSE/SPP1, May 2001, Department of Computer Science and Engineering, Indian Institute of Technology, Kharagpur, 721302, India.*, May 2001.
- [10] D. C. Prasad, S. P. Pal, and T. K. Dey. Visibility with multiple diffuse reflections. *Computational Geometry: Theory and Applications*, 10:187–196, 1998.
- [11] T. Shermer. Recent results in art galleries. *Proceeding of the IEEE*, 80:1384–1399, September 1992.